


TestroVax™

T-LEVEL BOOST*

A group of 43 men were divided into two groups: 1) the functional compound in TestroVax group and 2) an indistinguishable placebo group. All subjects consumed their test product every morning at breakfast for 12 consecutive days, and blood samples were taken before use, after 6 and 12 days of use, and 3 days after stopping use. All blood samples were drawn between 9:30-10:30 a.m., a time of day when serum oscillations of testosterone are at their minimum. Results showed that serum testosterone levels in the experimental group were significantly increased compared to those of the placebo group over the course of the study, wherein the key compound in TestroVax led to a 42.1% increase in

testosterone after 12 days of use. Interestingly, 3 days after the suspension of use, testosterone was still significantly increased compared to baseline in the experimental group, showing a lasting effect.*



Note to the People of California: The following notice is required by California's Safe Drinking Water and Toxic Enforcement Act of 1986, better known as Proposition 65. This notice does not apply to any other state in the Union.

WARNING: Consuming this product can expose you to lead, which is known to the State of California to cause birth defects or other reproductive harm. For more information, go to www.P65Warnings.ca.gov/food.

Formulated with a clinically validated compound to increase the natural production of testosterone and raise mean serum testosterone levels by 42.1%... **in just 12 days.***

Healthy testosterone levels have been associated with

- INTENSIFIED SEX DRIVE
- IMPROVED ENERGY & DRIVE
- MUSCLE MASS
- STRENGTH

OUR SATISFACTION GUARANTEE: Because no product or program will work for everyone, TestroVax™ is backed by our money-back guarantee. For details regarding our guarantee, please visit www.NovexBiotech.com or call customer service at: 1-800-898-5153.

This box was designed in an oversized manner to prevent pilferage. The actual size of the tablets is printed on the front of the box.

Manufactured for Novex Biotech®, LLC
Salt Lake City, UT 84116

Product resale allowed only through authorized representatives. International distribution is not authorized without prior written approval.

Store at controlled room temperature:
15° to 30°C / 59° to 86°F

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

©2021 ALL RIGHTS RESERVED. BR14082-10



DTPCFE A160MYFM.TI



TestroVax™

T-LEVEL BOOST*

CLINICALLY TESTED
TESTOSTERONE
BOOSTING
COMPOUND* **42.1%**
INCREASE
IN NATURAL
TESTOSTERONE
LEVELS* IN
JUST 12 DAYS!



DIETARY SUPPLEMENT

90 Tablets

Supplement Facts

Serving Size 3 Tablets
Servings Per Container 30

	Amount Per Serving	%Daily Value
Vitamin B6 (as pyridoxine hydrochloride)	2 mg	118%
Folate	667 mcg DFE (400 mcg folic acid)	167%
Vitamin B12 (as cyanocobalamin)	6 mcg	250%
Testrothione™ Proprietary Blend	2.7 g	
D-aspartic acid		†
Fenugreek (seed) Powder		†
L-arginine hydrochloride		†
Japanese Knotweed (root) Extract (containing resveratrol)		†
Grape (seed) Extract		†
Grape (skin) Extract		†
Azadirachtin (root) Extract		†
Licorice (root) Powder		†
Red Clover (aerial parts) Powder		†
Damiana (leaf) Extract		†
Ashwagandha (root) Extract		†
Taurine		†
Schizonepeta (aerial parts) Powder		†

†Daily Value not established.

Other ingredients: Microcrystalline cellulose, hydroxypropylcellulose, stearic acid, croscarmellose sodium, magnesium stearate, hydroxypropylmethylcellulose, polydextrose, titanium dioxide, silicon dioxide, talc, FD&C Blue 2 Lake, maltodextrin and medium chain triglycerides.

DIRECTIONS: Adult males take 3 tablets daily with breakfast. Individual results will vary.

WARNING: Do not exceed suggested daily serving. For adult males only. Before consuming TestroVax, consult your physician. Do not use TestroVax if you have any pre-existing medical condition, including but not limited to: high or low blood pressure, cardiac arrhythmia, stroke, heart, liver, or thyroid disease, anxiety, depression, seizure disorder, psychiatric disease, diabetes, pernicious anemia, difficulty urinating due to prostate enlargement, or if you are taking an MAO inhibitor or any other medication. Discontinue use if sleeplessness, tremors, dizziness, nervousness, headache, or heart palpitations occur. Do not use if the inner safety seal under the cap is broken or missing.

Use only as directed. Keep out of reach of children.

NOTICE: This product should not be consumed for more than 4 consecutive weeks with a 2- to 4-week washout period between cycles.

CAUTION: Increased testosterone levels may increase aggression, overconfidence, and sexual arousal.

Novex Biotech.

Novex Biotech.

Novex Biotech.